



قسم الأمراض الجلدية والتناسلية وطب الذكورة



اختبار أعمال السنة لطلبة السنة الخامسة طب
للعام الجامعي 2017/2016 م

المجموعة:

اسم الطالب:

الزمن المخصص: 15 دقيقة

التاريخ: 2017/ 1/ 24 م

1. Wheal:

- Is a secondary lesion.
- Varies in its color.
- Subsides with scar.
- Transient (1-2 hours).

2. Scutula are characteristic of infection with:

- Trichophyton violaceum.
- Trichophyton rubrum.
- Trichophyton schoenleinii.
- Microsporum audouinii.

3. Scabietic infestation is:

- Caused by notoedres.
- Self-limited disease.
- If treated, there is absolute immunity to scabies.
- If treated, the disease can be re-contracted again and again.

4. The variety of psoriasis that carry better prognosis is:

- Guttate.
- Erythrodermic.
- Pustular.
- Arthropathic.

5. Lesions of vitiligo:

- a. Occur in 4-6% of the world's population.
- b. Are asymptomatic.
- c. Do not fuse with neighboring lesions.
- d. Do not affect the pigment in associated hairs.

6. The course of a patch of alopecia areata is:

- a. Stationary.
- b. Gradual extension.
- c. Unpredictable.
- d. Appearance of further patches simultaneously in any part of the scalp.

7. The primary lesion of acne vulgaris is:

- a. Papule.
- b. Pustule.
- c. Nodule.
- d. Comedo.

8. The chancre of primary syphilis (untreated):

- a. Usually appears before the serology is positive.
- b. Always vanishes before secondary lesions appear.
- c. Is always single.
- d. Is usually tender.

9. All of the following statements about gonorrhoea in the adult females are true except:

- a. The condition is asymptomatic in 50% of cases.
- b. Vulva and vagina are sites of infection.
- c. Metastatic complications are similar to those in males.
- d. Urethra and cervix are sites of infection.

10. Hypospermia is diagnosed when the semen picture shows:

- a. Semen volume less than 2ml.
- b. Sperm count less than 20 million/ml.
- c. Sperm motility less than 20%.
- d. Abnormal Sperm morphology more than 20%.

Good Luck