



اختبار أعمال السنة لطلبة السنة الخامسة طب للعام الجامعي 2019/2018م

المجموعة:

اسم الطالب:

الزمن المخصص: 20 دقيقة

التاريخ : 30 / 10 / 2018 م

1. The following skin lesions could be considered as secondary lesions which may develop from modification of primary lesions except:

- a) Crust.
- b) Excoriations.
- c) Blister.
- d) Scales.

2. As regards tuberculoid leprosy, which is considered correct?

- a) Cutaneous lesions are many with symmetrical distribution.
- b) Nerve involvement is usually marked and in a few nerves.
- c) There are usually many bacilli in slit-skin smears.
- d) Lepromin test is usually weak.

3. 65 years old patient complained of severe pain in the trunk. At examination, grouped vesicles on an erythematous base were found. The most likely diagnosis will be:

- a) Bullous impetigo.
- b) Herpes Zoster.
- c) Chicken pox.
- d) Ramsay-Hunt syndrome.

4. Scutula is characteristic of infection with:

- a) Trichophyton violaceum.
- b) Trichophyton rubrum.
- c) Trichophyton schoenleinii.
- d) Microsporum audouinii.

5. Lichen planus is characterized by the following features except:

- a) Herald patch.
- b) Koebner phenomenon.
- c) Mucosal involvement.
- d) Violaceous flat topped polygonal papules.

6. Causes of non-cicatricial alopecia could include the following except:

- a) Alopecia areata.
- b) Trichotillomania.
- c) Kerion.
- d) Secondary syphilis.

7. One of the following features is correct as regards vitiligo:

- a) The autoimmune etiology is the only cause.
- b) The use of grafting techniques is possible in some cases.
- c) High association with hepatitis C virus.
- d) All of the above.

8. All of the following statements about gonorrhoea in the adult females are true except:

- a) The condition is asymptomatic in 50% of cases.
- b) Vulva and vagina are sites of infection.
- c) Metastatic complications are similar to those in males.
- d) Urethra and cervix are sites of infection.

9. The semen analysis of male infertile patient included semen volume 2.5 ml, sperm concentration 25×10^6 /ml, total sperm motility 60%, sperm abnormal forms 30% with pus cells 0.5 million/ml. your final diagnosis of the semen report will be:

- a) Astheno- teratozoospermia.
- b) Teratozoospermia with pyospermia.
- c) Teratozoospermia.
- d) Normal semen analysis.

10. The following are considered as oral lines of treatment of erectile dysfunction except:

- a) Yohimbine.
- b) Apomorphine.
- c) Prostaglandin E1.
- d) Sildenafil citrate.

Good Luck