
LYMPHATIC SYSTEM




LYMPH


■ What is lymph?

Tissue fluid (interstitial fluid) that enters the lymphatic vessels

FORMATION AND TRANSPORT OF TISSUE FLUID



LYMPHATIC SYSTEM



Essentially a drainage system accessory to venous system


larger particles that escape into tissue fluid can only be removed via lymphatic system

Functions of the Lymphatic System

- 24
- Reabsorbs excess interstitial fluid:
 - returns it to the venous circulation
 - maintain blood volume levels
 - prevent interstitial fluid levels from rising out of control.
- Transport dietary lipids:
 - transported through lacteals
 - drain into larger lymphatic vessels
 - eventually into the bloodstream.
- Iymphocyte development, and the immune response.


Components of the Lymphatic System

- 24-
 - Lymph
 - Lymphatic Vessels
 - **Lymphatic Capillaries**
 - Lymphatic Vessels
 - Lymphatic Trunks
 - Lymphatic Ducts
 - Lymphatic Organs
 - **Thymus**
 - Lymph Nodes
 - **□** Spleen
 - **□** Tonsils
 - Lymphatic cells



Lymph Vessels

- □ Lymphatic capillaries —
- Lymphatic collecting vessels
- □ Lymphatic trunks —
- □ Lymphatic ducts —




Lymphatic Capillaries

24

Features of structure:

- Blind end
- Single layer of overlapping endothelial cells
- More permeable than that of blood capillary
- Absent from avascular structures, brain, spinal cord splenic pulp and bone marrow



Lymphatic Capillaries – Lacteals


The small intestine contains special types of lymphatic capillaries called lacteals.

Lacteals pick up not only interstitial fluid, but also dietary lipids and lipid-soluble vitamins.


The lymph of this area has a milky color due to the lipid and is also called chyle. 2/-

Features of structure

- Three layered wall but thinner than vein,
- More numerous valves than in vein
- Interposed by lymph nodes at intervals
- Arranged in superficial and deep sets



LYMPH TRUNKS




LYMPHATIC DUCTS

Right lymphatic duct

 Formed by union of right jugular, subclavian, and bronchomediastinal trunks


Ends by entering the right venous angle



LYMPHATIC DUCTS


■Thoracic duct

- Begins in front of L1 as a dilated sac, the cisterna chyli
- formed by left and right lumbar trunks and intestinal trunk
- Enter thoracic cavity & ascends
- Travels upward, veering to the left at the level of T5



THORACIC DUCT.....

- At the root of the neck, it turns laterally
- arches forwards and descends to enter the left venous angle
- before termination, it receives the left jugular, Subclavian and bronchomediastinal trunk




DRAINAGE PATTERN

RIGHT LYMPHATIC DUCT -

Receives lymph from right half of head, neck, thorax and right upper limb, right lung, right side of heart, right surface of liver

THORACIC DUCT - Drains lymph from lower limbs, pelvic cavity, abdominal cavity, left side of thorax, and left side of the head, neck and left upper limb



Lymphatic Cells

- Also called lymphoid cells.
- Located in both the lymphatic system and the cardiovascular system.
- Work together to elicit an immune response.
- Types of lymphatic cells are:
 - macrophages
 - epithelial cells
 - dendritic cells
 - Iymphocytes

LYMPHATIC ORGANS

Primary organs


- Red bone marrow
- Thymus gland

Secondary organs

- Lymph nodes
- Lymph nodules
- ■Spleen

Lymph Nodes


- 24-
 - Small, round or oval
 - located along the pathways of lymph vessels.
 - length from 1 25 millimeters
 - Typically found in clusters
 - receive lymph from many body regions.
 - Lymph nodes are also found individually



Lymph node


Features

- Bean-shaped bodies
- With afferent vessels (entering at the periphery) and efferent lymph vessels(emerging at the hilus)
- Arranged in groups, along the blood vessels or the flexural side of the joint
- Divided into superficial and deep groups




Regional Lymph drainage

21



Regional Lymph Node is the lymph node where the lymph of the organ or part of the body drainge to firstly


Sentinel Lymph Node(in clinic)



Spleen


- Location
 - Left epigastric region
 - ≥ between 9th-11th rib
 - in line of 10th rib
- Largest lymphatic organ in the body.
- Can vary considerably in size and weight


■Function




Features

- Consists of two elongated lobes
- Is a large organ in the fetus
- Occupies the thoracic cavity behind the sternum
- Secrete lymphopoietin







Lymphatic Nodules

- Oval clusters of lymphatic cells with some extracellular matrix that are not surrounded by a connective tissue capsule.
- Filter and attack antigens.
- In some areas of the body, many lymphatic nodules group together to form larger structures.
 - mucosa-associated lymphatic tissue (MALT) or tonsils
 - very prominent in the mucosa of the small intestine, primarily in the ileum
 - Peyer patches
 - also present in the appendix


MALT


- MALT mucosa-associated lymphatic tissue:
 - Peyer's patches, tonsils, and the appendix (digestive tract)
 - Lymphoid nodules in the walls of the bronchi (respiratory tract)
- MALT protects the digestive and respiratory systems from foreign matter

Tonsils

- 24-
- clusters of lymphatic cells and extracellular matrix not completely surrounded by a connective tissue capsule.
- Consist of multiple germinal centers and crypts
- Several groups of tonsils form a protective ring around the pharynx.
 - pharyngeal tonsils (or adenoids) in nasopharynx
 - palatine tonsils in oral cavity
 - lingual tonsils along posterior one-third of the tongue


MALT (Mucosa Associated Lymphoid Tissue)






MALT (Peyer patches)







APPLIED ANATOMY

LYMPHANGITIS



Inflammation of the lymph vessels

Commonest cause bacteria called streptococcus pyogenes(most common).

Lymph vessels appear as red streaks through the skin



FILARIASIS

(b) Microfilaria develop into infective larvae in the mosquito and are injected into a new host

(c) Larvae mature into adult worms and spread through the lymphatic vessels, where they mate and lay eggs

LYMPHEDEMA

 Occurs due to accumulation of lymphatic fluid in the interstitial tissue

Sometimes can be appreciated after wearing tight clothing or jewellary on affected limb




LYMPHADENOPATHY

Means a disease of the lymph nodes

 Lymph nodes become swollen/ enlarged and may be painful to touch





LYMPHOMAS

 Cancers originating either from the lymphocytes in the lymph nodes or the lymphatic tissue in organs

Risk factors -- HIV,
HEPATITIS, EBV
infections



TONSILLITIS

- Infection of the pharyngeal tonsils
- □ Tonsils are swollen,
- Fever and pain during swallowing usually present
- Treatment surgical removal of tonsils (TONSILLECTOMY)




SPLENOMEGALY

- Enlarged Spleen
- Various causes



LET'S REVISE SOME IMPORTANT FACTS

Name the structures labelled A-E.



I am a part of lymphatic system and protect the body by clearing worn out red cells and foreign bodies from the blood stream. Who am I?



- A. Thymus
- **B.** Palatine Tonsil
- C. Spleen
- D. Lymph node

Which is the correct statement about the Lymphatic system?

- A. It Reabsorbs excess interstitial fluid and returns it to the venous circulation.
- B. Transports dietary lipids through lacteals.
- C. Helps in lymphocyte development, and the immune response.
- D. All of the above

Which disease caused by mosquit bite blocks the lymphatic drainage causing huge swelling of body parts (commonly lower lin

- A. Malaria
- **B.** Filaria
- C. Dengue
- D. Yellow fever



- Which of these is a primary lymphoid organ?
 - A. Lymph Node
 - B. Spleen
 - C. Tonsil
 - D. Bone Marrow