

Allied Academies Journals

Comparison of Difluprednate 0.05% versus Prednisolone Acetate 1% eye drops following uneventful cataract surgery --Manuscript Draft--

Manuscript Number:	Allied Academies-18-1116R1
Full Title:	Comparison of Difluprednate 0.05% versus Prednisolone Acetate 1% eye drops following uneventful cataract surgery
Short Title:	Difluprednate vs Prednisolone eyedrops post cataract surgery
Article Type:	Research Article
Section/Category:	Journal of Clinical Ophthalmology
Keywords:	Intraocular pressure, Difluprednate, Prednisolone acetate, steroid-induced ocular hypertension
Corresponding Author:	Islam El Saman, M.D Sohag, Egypt EGYPT
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	
Corresponding Author's Secondary Institution:	
First Author:	Islam El Saman, M.D
First Author Secondary Information:	
Order of Authors:	Islam El Saman, M.D Engy Mostafa, M.D Ahmed Gad Kamel, M.D Osama Ali, M.D
Order of Authors Secondary Information:	
Manuscript Region of Origin:	EGYPT
Abstract:	Purpose To compare intraocular pressure (IOP) readings and postoperative inflammation between 2 corticosteroid drops: difluprednate (DP) ophthalmic emulsion 0.05% and prednisolone acetate (PA) 1.0% after uneventful cataract surgery. Patients and methods randomized, controlled study of 343 patients undergoing uneventful cataract surgery in one eye. Randomization was used to allocate patients to either group: receive prednisolone acetate 1% five times daily or difluprednate 0.05% four times daily for 14 days post surgery, then tapered over 14 days. Intraocular pressure was measured preoperative, 1st day, 1st week, and one month postoperative. Also Ocular inflammation was monitored in the form of anterior chamber (AC) flare, AC cells, corneal edema and conjunctival congestion. Results No statistically significant difference in IOP between both groups was detected on postoperative first day and after one month. Yet there was statistically significant difference after first and second week of treatment with higher levels in the difluprednate group. As for ocular inflammation all signs resolved by the end of a

	month postoperative with no significant difference between both groups. Conclusion There was significant increased intraocular pressure in the diflurpednate group than in the prednisolone acetate group which might be attributed to the increased dosing and duration of use of difluprednate.
Suggested Reviewers:	
Opposed Reviewers:	
Response to Reviewers:	Revision made and IOP change written in percentage and significance was reported
Author Comments:	

Complete manuscript title: Comparison of Difluprednate 0.05% versus Prednisolone Acetate 1% eye drops following uneventful cataract surgery

- Running title: Difluprednate vs Prednisolone eye drops
post cataract surgery
- Full author name: Islam Saad El Saman MD, PhD*
Engy Mohamed Mostafa MD, PhD*
Ahmed Gad Kamel MD, PhD*
Osama Ali Mohammed MD, PhD*
- Affiliation: *Ophthalmology Department, Faculty of
Medicine, Sohag University, Sohag, 82524 Egypt
- Corresponding author: Isalm Saad El Saman
- Postal address: Ophthalmology Department, Sohag
University, Nasr City, Sohag, Egypt. 82524
- E-mail address: eslamsaad@med.sohag.edu.eg
- Telephone number: +2/01212391940
- Conflict of interest statement: The authors have no
proprietary interests or conflicts of interest related to this
submission.

- This submission has not been published anywhere previously and that it is not simultaneously being considered for any other publication.
- No author has any financial or proprietary interest in a product, method, or material mentioned in this manuscript
- Disclosure of any funding received for this work: NONE

Abstract

Purpose: To compare intraocular pressure (IOP) readings and postoperative inflammation between 2 corticosteroid drops: difluprednate (DP) ophthalmic emulsion 0.05% and prednisolone acetate (PA) 1.0% after uneventful cataract surgery.

Patients and methods: randomized, controlled study of 343 patients undergoing uneventful cataract surgery in one eye. Randomization was used to allocate patients to either group: receive prednisolone acetate 1% five times daily or difluprednate 0.05% four times daily for 14 days post surgery, then tapered over 14 days. Intraocular pressure was measured preoperative, 1st day, 1st week, and one month postoperative. Also, ocular inflammation was monitored in the form of anterior chamber (AC) flare, AC cells, corneal edema and conjunctival congestion.

Results: No statistically significant difference in IOP between both groups was detected on postoperative first day and after one month. Yet there was statistically significant difference after first and second week of treatment with higher levels in the difluprednate group. As for ocular inflammation all signs resolved by the end of a month postoperative with no significant difference between both groups.

Conclusion: There was significant increased intraocular pressure in the difluprednate group than in the prednisolone acetate group which might be attributed to the increased dosing and duration of use of difluprednate.

Keywords: Intraocular pressure, Difluprednate, Prednisolone acetate, steroid-induced ocular hypertension

Introduction

Topical ocular corticosteroids has been the main stay after intraocular surgery¹ as they suppress ocular inflammation and help improve visual outcomes.^{2, 3} Postoperative inflammation is a natural result of tissue injury and corticosteroids acts by inhibiting the production of both leukotrienes and prostaglandins.⁴

Despite their importance in management of inflammation and speeding up recovery, side effects are inevitable.^{5, 6} The most dreaded complication is increased intraocular pressure especially steroid responders who constitute 5% of population.⁷

Prednisolone acetate 1% (PA) being the vastly used topical corticosteroid therapy. Yet PA 1% usually necessitates multiple applications especially in severe cases;⁸ which might increase noncompliance, leading to under achievement of therapeutic goals.

Difluprednate ophthalmic emulsion 0.05% is a recently available topical ocular corticosteroid that was approved in 2008 initially for treatment of inflammation and pain that develops with ocular surgery, and for management of endogenous anterior uveitis.⁹ Difluprednate is a prednisolone acetate derivative¹⁰ that is strengthened by two fluorinations at carbons 6 and 9 thus enhancing its potency, a butyrate group at carbon 17 adding anti inflammatory activity, and an acetic acid group at carbon 21 to increase penetration.¹¹

Difluprednate 0.05% was reported to be effective at decreasing inflammation and pain in postoperative ocular surgery patients.^{12, 13}

This study aimed to detect whether difluprednate 0.05% dosed four times daily is as effective as prednisolone acetate 1% dosed five times daily in patients undergoing uneventful cataract surgery (phacoemulsification). In addition to monitor IOP after both eye drops postoperatively.

Patients and methods

This is a prospective randomized case controlled study of 343 patients undergoing phacoemulsification in Sohag University Hospital, Sohag, Egypt between January –June 2018. The study protocol was approved by the ethical committee of Sohag Faculty of Medicine with informed written consent being obtained from all patients. Exclusion criteria included: history of glaucoma or ocular hypertension in the study eye, previous history of steroid-related intraocular pressure (IOP) increase, or at the time of recruiting had an IOP > 24 mmHg in the operated eye; previous ocular surgery; history of uveitis; intraoperative complications as posterior capsule rupture and diabetic patients. Patients were subjected to routine ophthalmological examination: visual acuity assessment, slit lamp examination, ICare tonometer was used for intraocular pressure measurement after being calibrated.

Patients who met inclusion criteria were randomized in a 1:1 ratio to receive either Prednisolone acetate or difluprednate. All patients underwent phacoemulsification by a single surgeon (OA) with intraocular lens implantation. The 2 eye drops were initiated after surgery and continued for one month. The prescribed dose of PA was one drop 5 times daily for 15 days then tapered to twice daily on 3rd week then once on the 4th week. While DP was administered 4 times daily for 15 days then

tapered to twice daily on 3rd week then once on the 4th week.

Patients were instructed for self-administration.

During follow up period, doses and method of administration of steroids drops were adjusted if the treating physician detected that treatment response was inadequate or hazardous at any time point.

The primary outcome was the incidence of an increase in IOP higher than 6 mm Hg from the preoperative IOP or IOP higher than 21 mm Hg 1 month postoperatively.

Ocular inflammation assessment included AC flare, AC cell count, bulbar conjunctival injection or ciliary injection, and corneal edema. Ocular inflammation grading criteria is listed in table 1. Control of ocular inflammation was achieved when AC cell grade of 0 (≤ 5 cells) and a flare grade of 0 (complete absence) were noted on day 14 postoperative.

Follow-up visits were every week for 4 weeks. The study was in compliance with the tenets of the Helsinki's Declaration, and informed consent was obtained from all study patients.

Table 1. ¹⁴ Ocular inflammation diagnostic criteria	
AC cell count	Grade 0 ≤ 5 cells Grade 1 6–15 cells Grade 2 16–25 cells Grade 3 26–50 cells Grade 4 > 50 cells
AC flare	Grade 0 Complete absence Grade 1 Very slight

	Grade 2 Moderate Grade 3 Marked Grade 4 Intense
Bulbar conjunctival injection, ciliary injection, corneal edema	Grade 0 Absent Grade 1 Mild Grade 2 Moderate Grade 3 Severe

Statistical analysis

Descriptive statistics was used for characteristics of both groups. Differences between the 2 groups were assessed with sample t tests and chi-square or Fisher exact tests for person-based measures. Statistical Analysis Software version (SPSS) 9.4 was used for all statistical analyses.


Results

This prospective randomized study included 343 patients (178 males and 165 females) with a mean age of 48 ± 7.3 years (range: 42-65). The PA group included 169 eyes while DP group included 174 eyes. There was no statistically significant difference between both groups as regards age, sex and preoperative IOP. (Table 2)

Table 2. Demographics of PA and DP group			
	Prednisolone acetate group (n=169)	Difluprednate group (n=174)	P value

Age (yrs)	47±6.6	48±5.3	0.09
Mean±SD (range)	(42-62)	(41-65)	
Sex (M/F)	89/80	89/85	
Preoperative IOP (mmHg)	12±3.6	13±3.1	0.23
Mean±SD (range)	(11-16)	(11-15)	

Table 3. IOP measurement over the follow up period			
	Prednisolone acetate group (n=169)	Difluprednate group (n=174)	P value
Postoperative 1 st day	14.2±3.8	14.5±5.1	0.33
Postoperative 1 st week	15.8±4.5	16.5±5.5	0.03
Postoperative 2nd week	15.2±4.6	15.7±5.4	0.043
Postoperative 3rd week	14.9±4.8	15.1±5	0.094
Postoperative one month	14.6±3.9	14.9±4.4	0.26



There was no statistically significant difference in IOP between both groups on postoperative first day and after one month. Yet

there was statistically significant difference after first and second week of treatment (Table 3). The total number of patients in PA and DP group who showed significant change in IOP was: 7 (4.14%) and 14 (8%) patients respectively which was statistically significantly different. On 1st week follow up of both groups, 4 eyes (2.4%) in the PA group and 7 eyes (4%) in the DP group showed a 6mmHg increase in IOP higher than preoperatively. The IOP increase percentage in the first week in the PA and DP group was 10% and 12% respectively which showed significant difference ($p=0.05$). On the other hand, 2 eyes (1.2%) in the PA group and 5 eyes (2.9%) in the DP group had an IOP increase higher than 21 mm Hg. In cases of increase IOP 26 mmHg and below, the steroid eye drops were stopped and substituted with NSAID eye drops twice daily and followed day after day until stabilization of IOP. Three eyes developed increase IOP >30mmHg; one eye in PA group and 2 in DP group: anti-glaucoma eye drop was added (dorzolamide and timolol 0.5% combination) twice daily until IOP was normalized. None of these patients developed permanent glaucomatous optic neuropathy.

As for the ocular inflammation follow up data, it is represented in table 4 with no statistical difference at the end of the follow up month up between both groups. Yet the resolution of signs of ocular inflammation was faster in the PA group after one week of treatment.

Table 4. Postoperative ocular inflammation follow up							
		Postop 1 st day		Postop 1 st week		Postop 4 th week	
		PA group	DP group	PA group	DP group	PA group	DP group
AC cell count	Grade 0	130	125	160	160	169	173
	Grade 1	33	40	9	12	0	1
	Grade 2	6	9	0	2	0	0
	Grade 3	0	0	0	0	0	0
	Grade 4	0	0	0	0	0	0
AC flare	Grade 0	133	109	155	152	169	170
	Grade 1	28	50	14	20	0	4
	Grade 2	8	11	0	2	0	0
	Grade 3	0	0	0	0	0	0
	Grade 4	0	0	0	0	0	0
Bulbar conjunctival injection, ciliary injection, corneal edema	Grade 0	140	139	155	154	169	171
	Grade 1	26	30	14	20	0	3
	Grade 2	3	5	0	0	0	0
	Grade 3	0	0	0	0	0	0

Discussion

In our comparative study, we aimed to report corticosteroid induced glaucoma and persistence of post operative inflammation after the use of one of two eye drops which are: the most commonly used Prednisolone Acetate 1% and the newer Difluprednate 0.05% eye drop. Difluprednate ophthalmic emulsion 0.05% (DFBA) (Diflustero, Orchidia, Egypt) is a recent medication used in the treatment of postoperative inflammation.¹⁵ It was FDA approved in July 2008 for the treatment of inflammation and pain post ocular surgery. DFBA is different from PA in its chemical structure which allows 56 times increased receptor-binding affinity to glucocorticoid

receptors,¹⁰ resulting in increased anti-inflammatory capacity, and increased tissue penetration.¹¹

Corticosteroid-induced ocular hypertension has multiple definitions. Becker¹⁶ used an IOP of 31mmHg as the cutoff value for stopping the use of betamethasone. While Armaly¹⁷ classified IOP steroid response increase into 3 groups: low (≤ 5 mm Hg), intermediate (6 to 15 mm Hg), and high (≥ 16 mm Hg) with 6 mm Hg as the lower limit of a clinically significant response. On the other hand, Stewart et al¹⁸ proposed a significant would be ≥ 10 mm Hg. In our study difluprednate group showed greater number of eyes with increased IOP either 6mmHg above the preoperative pressure or even eyes above 30 mmHg. In 2014, Jeng et al.¹⁹ carried out a retrospective review of treatments after vitreoretinal surgeries and found that eyes that received difluprednate were at an higher risk of developing a clinically significant increase in IOP than eyes receiving with prednisolone acetate.

Patients who have an excessive amount of postoperative inflammation would benefit more from difluprednate use.²⁰ In 1990, Eisen et al²¹ reported that less complicated dosing regimens result in better adherence. Yet the difluprednate group did not show superior results to the prednisolone group. In ophthalmologic clinical practice, DFBA has been shown to be effective in reducing postoperative inflammation and decreasing corneal edema and retinal thickness.^{13-15, 22} However, reports

have shown an increase in IOP when used after ophthalmologic surgery.^{19, 23, 24}

There is an ongoing debate as to what is the most effective postoperative treatment after cataract surgery. Kessel et al²⁵ compared the use of non steroidal anti-inflammatory drugs (NSAIDs) with the use of topical steroids in patients who had uneventful cataract surgery. They reported that topical NSAIDs were more effective in decreasing inflammation and postoperative cystoid macular edema with no differences in side events or visual results.

In addition, difluprednate was shown to have increased bioavailability and dose uniformity owing to the emulsion form rather than a suspension.¹² So the main benefit of difluprednate, would be frequent dosing with superior dose uniformity.¹³

Our results showed that there was a significantly greater incidence of corticosteroid-induced IOP response in eyes receiving difluprednate than eyes treated with prednisolone acetate for postoperative inflammation after cataract surgery.

References:

- [1] Optometric Clinical Practice Guideline: Care of the Patient With Anterior Uveitis. St. Louis, MO: American Optometric Association. , 1994.
- [2] Lorenz K, Dick B, Jehkul A, Auffahrt GU. Inflammatory response after phacoemulsification treated with 0.5%

prednisolone acetate or vehicle. Graefes Arch Clin Exp Ophthalmol 2008;246(11):1617-22.

[3] Rajpal RK, Fong R, Comstock TL. Loteprednol etabonate ophthalmic gel 0.5% following cataract surgery: integrated analysis of two clinical studies. Adv Ther 2013;30(10):907-23.

[4] Rugstad HE. Antiinflammatory and immunoregulatory effects of glucocorticoids: mode of action. Scand J Rheumatol Suppl 1988;76:257-64.

[5] Carnahan MC, Goldstein DA. Ocular complications of topical, peri-ocular, and systemic corticosteroids. Curr Opin Ophthalmol 2000;11(6):478-83.

[6] Kersey JP, Broadway DC. Corticosteroid-induced glaucoma: a review of the literature. Eye (Lond) 2006;20(4):407-16.

[7] Razeghinejad MR, Katz LJ. Steroid-induced iatrogenic glaucoma. Ophthalmic Res 2012;47(2):66-80.

[8] Vandenbroeck S, De Geest S, Dobbels F, et al. Prevalence and correlates of self-reported nonadherence with eye drop treatment: the Belgian Compliance Study in Ophthalmology (BCSO). J Glaucoma 2011;20(7):414-21.

[9] Sheppard JD, Toyos MM, Kempen JH, et al. Difluprednate 0.05% versus prednisolone acetate 1% for endogenous anterior uveitis: a phase III, multicenter, randomized study. Invest Ophthalmol Vis Sci 2014;55(5):2993-3002.

[10] Tajika T, Waki M, Tsuzuki M, et al. Pharmacokinetic features of difluprednate ophthalmic emulsion in rabbits as determined by glucocorticoid receptor-binding bioassay. J Ocul Pharmacol Ther 2011;27(1):29-34.

[11] Bodor N, Harget AJ, Phillips EW. Structure-activity relationships in the antiinflammatory steroids: a pattern-recognition approach. J Med Chem 1983;26(3):318-28.

[12] Donnenfeld ED, Holland EJ, Solomon KD, et al. A multicenter randomized controlled fellow eye trial of pulse-dosed difluprednate 0.05% versus prednisolone acetate 1% in cataract surgery. Am J Ophthalmol 2011;152(4):609-17 e1.

[13] Korenfeld MS, Silverstein SM, Cooke DL, et al. Difluprednate ophthalmic emulsion 0.05% for postoperative

inflammation and pain. J Cataract Refract Surg 2009;35(1):26-34.

[14] Smith S, Lorenz D, Peace J, et al. Difluprednate ophthalmic emulsion 0.05% (Durezol) administered two times daily for managing ocular inflammation and pain following cataract surgery. Clin Ophthalmol 2010;4:983-91.

[15] Jamal KN, Callanan DG. The role of difluprednate ophthalmic emulsion in clinical practice. Clin Ophthalmol 2009;3:381-90.

[16] Becker B. Intraocular Pressure Response to Topical Corticosteroids. Invest Ophthalmol 1965;4:198-205.

[17] Armaly MF. Statistical Attributes of the Steroid Hypertensive Response in the Clinically Normal Eye. I. The Demonstration of Three Levels of Response. Invest Ophthalmol 1965;4:187-97.

[18] Stewart RH, Smith JP, Rosenthal AL. Ocular pressure response to fluorometholone acetate and dexamethasone sodium phosphate. Curr Eye Res 1984;3(6):835-9.

[19] Jeng KW, Fine HF, Wheatley HM, et al. Incidence of steroid-induced ocular hypertension after vitreoretinal surgery with difluprednate versus prednisolone acetate. Retina 2014;34(10):1990-6.

[20] Stringer W, Bryant R. Dose uniformity of topical corticosteroid preparations: difluprednate ophthalmic emulsion 0.05% versus branded and generic prednisolone acetate ophthalmic suspension 1%. Clin Ophthalmol 2010;4:1119-24.

[21] Eisen SA, Miller DK, Woodward RS, et al. The effect of prescribed daily dose frequency on patient medication compliance. Arch Intern Med 1990;150(9):1881-4.

[22] Foster CS, Davanzo R, Flynn TE, et al. Durezol (Difluprednate Ophthalmic Emulsion 0.05%) compared with Pred Forte 1% ophthalmic suspension in the treatment of endogenous anterior uveitis. J Ocul Pharmacol Ther 2010;26(5):475-83.

[23] Meehan K, Vollmer L, Sowka J. Intraocular pressure elevation from topical difluprednate use. Optometry 2010;81(12):658-62.

- [24] Birnbaum AD, Jiang Y, Tessler HH, Goldstein DA. Elevation of intraocular pressure in patients with uveitis treated with topical difluprednate. *Arch Ophthalmol* 2011;129(5):667-8.
- [25] Kessel L, Tendal B, Jorgensen KJ, et al. Post-cataract prevention of inflammation and macular edema by steroid and nonsteroidal anti-inflammatory eye drops: a systematic review. *Ophthalmology* 2014;121(10):1915-24.